

Rewriting Preserving Recognizability of Finite Tree Languages

Sándor Vágvolgyi*

*Department of Foundations of Computer Science
University of Szeged
Szeged, Árpád tér 2
H-6720 Hungary*

Abstract

We show that left-linear generalized semi-monadic TRSs effectively preserving recognizability of finite tree languages (are EPRF-TRSs). We show that reachability, joinability, and local confluence are decidable for EPRF-TRSs.

Keywords: term rewrite systems, tree automata, preservation of recognizability

1 Introduction

The notion of preservation of recognizability through rewriting is a widely studied concept in term rewriting, see [2]-[5], [7], [10]-[22]. Let Σ be a ranked alphabet, let R be a term rewrite system (TRS) over Σ , and let L be a tree language over Σ . Then $R_\Sigma^*(L)$ denotes the set of descendants of trees in L . A TRS R over Σ preserves Σ -recognizability (is a P Σ R-TRS), if for each recognizable tree language L over Σ , $R_\Sigma^*(L)$ is recognizable. A TRS R over Σ preserves Σ -recognizability of finite tree languages (is a P Σ RF-TRS), if for each finite tree language L over Σ , $R_\Sigma^*(L)$ is recognizable.

Let R be a TRS over Σ . Then its signature, $\text{sign}(R) \subseteq \Sigma$ is the ranked alphabet consisting of all symbols appearing in the rules of R . A TRS R over $\text{sign}(R)$ preserves recognizability (is a PR-TRS), if for each ranked alphabet Σ with $\text{sign}(R) \subseteq \Sigma$, R , as a TRS over Σ , preserves Σ -recognizability. A TRS R over $\text{sign}(R)$ preserves recognizability of finite tree languages (is a PRF-TRS), if for each ranked alphabet Σ with $\text{sign}(R) \subseteq \Sigma$, R , as a TRS over Σ , preserves Σ -recognizability of finite tree languages.

*E-mail: vagvolgy@inf.u-szeged.hu.

A TRS R over Σ effectively preserves Σ -recognizability (is an EP Σ R-TRS), if for a given a bottom-up tree automaton (bta) $\mathcal{B}$ over Σ , we can effectively construct a bta $\mathcal{C}$ over Σ such that $L(\mathcal{C}) = R_{\Sigma}^*(L(\mathcal{B}))$. A TRS R over Σ effectively preserves Σ -recognizability of finite tree languages (is an EP Σ RF-TRS), if for a given finite tree language L over Σ , we can effectively construct a bta $\mathcal{C}$ over Σ such that $L(\mathcal{C}) = R_{\Sigma}^*(L)$. A TRS R over $sign(R)$ effectively preserves recognizability of finite tree languages (is an EPRF-TRS), if for a given ranked alphabet Σ with $sign(R) \subseteq \Sigma$ and a given finite tree language L over Σ , we can effectively construct a bta $\mathcal{C}$ over Σ such that $L(\mathcal{C}) = R_{\Sigma}^*(L(\mathcal{B}))$.

Gyenizse and Vágvolgyi [12] presented a linear TRS R over $sign(R)$ such that R is an EP $sign(R)$ R-TRS and R is not a PR-TRS. Vágvolgyi [22] showed that it is not decidable for a murg TRS R over Σ whether R is a P Σ RF-TRS. Let R be a rewrite system over $sign(R)$, and let $\Sigma = \{f, \sharp\} \cup sign(R)$, where $f \in \Sigma_2 - sign(R)$ and $\sharp \in \Sigma_0 - sign(R)$. Gyenizse and Vágvolgyi [12] showed that R is an EP Σ R-TRS if and only if R is an EPR-TRS. Gyenizse and Vágvolgyi [13] improved this result for left-linear TRSs. They showed the following. Let R be a left-linear TRS over $sign(R)$, and let $\Sigma = \{g, \sharp\} \cup sign(R)$, where $g \in \Sigma_1 - sign(R)$ and $\sharp \in \Sigma_0 - sign(R)$. Then R is an EP Σ R-TRS if and only if R is an EPR-TRS.

In [10] Gilleron showed that for a TRS R over Σ it is not decidable whether R is a P Σ R-TRS. We may naturally introduce the above concepts for string rewrite systems as well. Otto [16] has proved that a string rewrite system R over the alphabet $alph(R)$ of R preserves $alph(R)$ -recognizability if and only if R preserves recognizability. Otto [16] showed that it is not decidable for a string rewrite system R whether R preserves $alph(R)$ -recognizability, and whether R preserves recognizability. Hence it is not decidable for a linear TRS R whether R is a PR-TRS, see [16].

In spite of the undecidability results of Gilleron [10] and Otto [16], we know several classes of EPR-TRSs. Gyenizse and Vágvolgyi [12] generalized the concept of a semi-monadic TRS [2] introducing the concept of a generalized semi-monadic TRS (GSM-TRS for short). They showed that each linear GSM-TRS R is an EPR-TRS. Takai, Kaji, and Seki [18] introduced finite path overlapping TRS's (FPO-TRSs). They [18] showed that each right-linear FPO-TRS R is an EPR-TRS. They [18] also showed that each GSM-TRS R is an FPO-TRS. Thus we get that that each right-linear GSM-TRS R is an EPR-TRS. Vágvolgyi [20] introduced the concept of a half-monadic TRS. Each right-linear half-monadic TRS is an FPO-TRS. Hence each right-linear half-monadic TRS is an EPR-TRS. Using this result, Vágvolgyi [20] showed that termination and convergence are decidable properties for right-linear half-monadic term rewrite systems. Takai, Seki, Fujinaka, and Kaji [19] presented an EPR-TRS which is not an FPO-TRS, see Example 1 in [19]. Takai, Seki, Fujinaka, and Kaji [19] introduced layered transducing term rewriting systems (LT-TRSs). They [19] showed that each right-linear LT-TRS R is an EPR-TRS.

We show that each terminating TRS is an EPRF-TRS. We adopt the construction of Salomaa [17], Coquidé et al [2], and Gyenizse and Vágvolgyi [12], when showing that any left-linear GSM-TRS R is an EPRF-TRS. We slightly modify the

proofs of the decision results of Gyenizse and Vágvolgyi [12] when we show the following decidability results.

(1) Let R be an EPRF-TRS over Σ , and let $p, q \in T_\Sigma(X)$. Then it is decidable whether $p \rightarrow_R^* q$.

(2) Let R be an EPRF-TRS over Σ , and let $p, q \in T_\Sigma(X)$. Then it is decidable whether there exists a tree $r \in T_\Sigma(X)$ such that $p \rightarrow_R^* r$ and $q \rightarrow_R^* r$.

(3) Let R be a confluent EPRF-TRS over Σ , and let $p, q \in T_\Sigma(X)$. Then it is decidable whether $p \leftrightarrow_R^* q$.

(4) For an EPRF-TRS R , it is decidable whether R is locally confluent.

(5) Let R be an EPRF-TRS, and let S be a TRS over Σ . Then it is decidable whether $\rightarrow_S^* \subseteq \rightarrow_R^*$.

(6) Let R and S be EPRF-TRSs. Then it is decidable which one of the following four mutually excluding conditions hold.

- (i) $\rightarrow_R^* \subset \rightarrow_S^*$,
- (ii) $\rightarrow_S^* \subset \rightarrow_R^*$,
- (iii) $\rightarrow_R^* = \rightarrow_S^*$,
- (iv) $\rightarrow_R^* \bowtie \rightarrow_S^*$,

where “ $\bowtie$ ” stands for the incomparability relationship.

(7) Let R be an EPRF-TRS. Then it is decidable whether R is left-to-right minimal. (A TRS R is left-to-right minimal if for each rule $l \rightarrow r$ in R , $\rightarrow_{R-\{l \rightarrow r\}}^* \subset \rightarrow_R^*$.)

(8) Let R and S be TRSs such that $R \cup R^{-1}$ and $S \cup S^{-1}$ are EPRF-TRSs. Then it is decidable which one of the following four mutually excluding conditions holds.

- (i) $\leftrightarrow_R^* \subset \leftrightarrow_S^*$,
- (ii) $\leftrightarrow_S^* \subset \leftrightarrow_R^*$,
- (iii) $\leftrightarrow_R^* = \leftrightarrow_S^*$,
- (iv) $\leftrightarrow_R^* \bowtie \leftrightarrow_S^*$.

Fülöp’s [5] undecidability results on deterministic top-down tree transducers simply imply the following. Each of the following questions is undecidable for any convergent left-linear EPRF-TRSs R and S over a ranked alphabet Ω , for any recognizable tree language $L \subseteq T_\Omega$ given by a tree automaton over Ω recognizing L . Here $\Gamma \subseteq \Omega$ is the smallest ranked alphabet for which $R(L) \subseteq T_\Gamma$. Furthermore, the set of R -normal forms of the trees in L is denoted by $R(L)$.

- (i) Is $R(L) \cap S(L)$ empty?
- (ii) Is $R(L) \cap S(L)$ infinite?
- (iii) Is $R(L) \cap S(L)$ recognizable?
- (iv) Is $T_\Gamma - R(L)$ empty?
- (v) Is $T_\Gamma - R(L)$ infinite?
- (vi) Is $T_\Gamma - R(L)$ recognizable?
- (vii) Is $R(L)$ recognizable?
- (viii) Is $R(L) = S(L)$?
- (ix) Is $R(L) \subseteq S(L)$?

Fülöp and Gyenizse [6] showed that it is undecidable for a tree function induced by a deterministic homomorphism whether it is injective. Hence for any conver-

gent left-linear EPRF-TRS R over a ranked alphabet Σ , and any recognizable tree language $L \subseteq T_\Sigma$, it is undecidable whether the tree function $\rightarrow_R^* \cap (L \times R(L))$ is injective.

Finally we show the following. Let R be a linear collapse-free EPRF-TRS and S be a linear collapse-free EPR-TRS over the disjoint ranked alphabets $\text{sign}(R)$ and $\text{sign}(S)$, respectively. Then $R \oplus S$ is a linear collapse-free EPR-TRS.

This paper is divided into seven sections. In Section 2, we recall the necessary notions and notations. In Section 3, we study TRS classes preserving recognizability. In Section 4, we show that left-linear GSM-TRSs are EPRF-TRSs. In Section 5, we illustrate the constructions presented in Section 4 by an example. In Section 6, we show various decidability and undecidability results on PRF-TRSs and EPRF-TRSs. Finally, in Section 7, we present our concluding remarks, and some open problems.

2 Preliminaries

We recall and invent some notations, basic definitions and terminology which will be used in the rest of the paper. Nevertheless the reader is assumed to be familiar with the basic concepts of term rewrite systems and of tree language theory (see, e.g. Baader, Nipkow[1], Gécseg, Steinby [8], [9]).

2.1 Terms

The cardinality of a set A is denoted by $|A|$. We denote by ρ^{-1} the inverse of a binary relation ρ . The composition of binary relations ρ and τ is denoted by $\rho \circ \tau$.

The set of nonnegative integers is denoted by N , and N^* stands for the free monoid generated by N with empty word λ as identity element. For a word $\alpha \in N^*$, $\text{length}(\alpha)$ stands for the length of α .

A ranked alphabet is a finite set Σ in which every symbol has a unique rank in N . For $m \geq 0$, Σ_m denotes the set of all elements of Σ which have rank m . The elements of Σ_0 are called constants. We assume that all ranked alphabets Σ and Δ that we consider have the following property. If $\sigma \in \Sigma_i$, and $\sigma \in \Delta_j$, then $i = j$. In other words, σ has the same rank in Σ as in Δ .

For a set of variables Y and a ranked alphabet Σ , $T_\Sigma(Y)$ denotes the set of Σ -terms (or Σ -trees) over Y . $T_\Sigma(\emptyset)$ is written as T_Σ . A term $t \in T_\Sigma$ is called a ground term. A tree $t \in T_\Sigma(Y)$ is linear if any variable of Y occurs at most once in t . We specify a countable set $X = \{x_1, x_2, \dots\}$ of variables which will be kept fixed in this paper. Moreover, we put $X_m = \{x_1, \dots, x_m\}$, for $m \geq 0$. Hence $X_0 = \emptyset$.

For any $m \geq 0$, we distinguish a subset $\bar{T}_\Sigma(X_m)$ of $T_\Sigma(X_m)$ as follows: a tree $t \in T_\Sigma(X_m)$ is in $\bar{T}_\Sigma(X_m)$ if and only if each variable in X_m appears exactly once in t .

For a term $t \in T_\Sigma(X)$, the height $\text{height}(t) \in N$, the set of variables $\text{var}(t)$ of t , and the set of positions $\text{POS}(t) \subseteq N^*$ are defined in the usual way. For each $t \in T_\Sigma(X)$ and $\alpha \in \text{POS}(t)$, $t/\alpha \in T_\Sigma(X)$ is the subterm of t at α . For a tree $t \in$

$T_\Sigma(X)$, $sub(t)$ denotes the *subtrees* of t . For a tree language $L \subseteq T_\Sigma$, the set $sub(L)$ of subtrees of elements of L is defined by the equality $sub(L) = \bigcup (sub(t) \mid t \in L)$.

For $t \in T_\Sigma$, $\alpha \in POS(t)$, and $r \in T_\Sigma$, we define $t[\alpha \leftarrow r] \in T_\Sigma$ as follows.

- (i) If $\alpha = \lambda$, then $t[\alpha \leftarrow r] = r$.
- (ii) If $\alpha = i\beta$, for some $i \in N$ and $\beta \in N^*$, then $t = f(t_1, \dots, t_m)$ with $f \in \Sigma_m$ and $1 \leq i \leq m$. Then $t[\alpha \leftarrow r] = f(t_1, \dots, t_{i-1}, t_i[\beta \leftarrow r], t_{i+1}, \dots, t_m)$.

Let Σ be a ranked alphabet. Let $f \in \Sigma_1$, $t \in T_\Sigma$ be arbitrary. The tree $f^k(t) \in T_\Sigma$, $k \geq 0$, is defined by recursion: $f^0(t) = t$, and $f^{k+1}(t) = f(f^k(t))$ for $k \geq 0$.

A substitution is a mapping $\sigma : X \rightarrow T_\Sigma(X)$ which is different from the identity only for a finite subset $Dom(\sigma)$ of X . For any substitution σ with $Dom(\sigma) \subseteq X_m$, $m \geq 0$, the term $\sigma(t)$ is produced from t by replacing each occurrence of x_i with $\sigma(x_i)$ for $1 \leq i \leq m$. For any trees $t \in \bar{T}(X_k)$, $t_1, \dots, t_k \in T_\Sigma(X)$ and for the substitution σ with $Dom(\sigma) \subseteq X_k$ and $\sigma(x_i) = t_i$ for $i = 1, \dots, k$, we denote the term $\sigma(t)$ by $t[t_1, \dots, t_k]$ as well. Moreover, for any k, m with $1 \leq m \leq k$, for any tree $t \in T_\Sigma(\{x_m, \dots, x_k\})$ and for any substitution σ with $\sigma(x_m) = t_m, \dots, \sigma(x_k) = t_k$, we denote $\sigma(t)$ also by $t[x_m \leftarrow t_m, \dots, x_k \leftarrow t_k]$.

Let Σ be a ranked alphabet and let $u, v \in T_\Sigma(X)$. The tree u is a supertree of v if u is linear and there is a substitution σ such that $v = \sigma(u)$.

We say that the pair (l_1, r_1) is a variant of the pair (l_2, r_2) if there is a substitution $\sigma : X \rightarrow X$ such that

- (i) $\sigma(l_2) = l_1$ and $\sigma(r_2) = r_1$, and that
- (ii) for all $x_i, x_j \in var(l_2) \cup var(r_2)$, $\sigma(x_i) = \sigma(x_j)$ implies that $x_i = x_j$.

For the concept of a unifier and a most general unifier (mgu), see [1].

2.2 TRSs

Let Σ be a ranked alphabet. Then a term rewrite system (TRS) R over Σ is a finite subset of $T_\Sigma(X) \times T_\Sigma(X)$ such that for each $(l, r) \in R$, each variable of r also occurs in l . Elements (l, r) of R are called rules and are denoted by $l \rightarrow r$. Furthermore, $sign(R) \subseteq \Sigma$ is the ranked alphabet consisting of all symbols appearing in the rules of R .

Let R be a TRS over Σ . Given any two terms s and t in $T_\Sigma(X)$ and an position $\alpha \in POS(s)$, we say that s rewrites to t at α and denote this by $s \rightarrow_R t$ if there is some pair $(l, r) \in R$ and a substitution σ such that $s/\alpha = \sigma(l)$ and $t = s[\alpha \leftarrow \sigma(r)]$. Here we also say that R rewrites s to t applying the rule $l \rightarrow r$ at α .

Reachability, joinability, termination, confluence, local confluence, convergence are defined in the usual way, see [1].

We say that a TRS R is collapse-free if there is no rule $l \rightarrow r$ in R such that $l \in X$ or $r \in X$.

A left-linear (linear, resp.) TRS is one in which no variable occurs more than once on any left-hand side (right-hand side and left-hand side, resp.). A ground TRS is one of which all rules are ground (i.e. elements of $T_\Sigma \times T_\Sigma$).

A TRS is monadic if each left-hand side is of height at least 1 and each right-hand side is of height at most 1. A TRS is called right-ground if each right-hand side is ground. A TRS R over Σ is *murg* if R is the union of a monadic TRS and a right-ground TRS over Σ . Obviously, each monadic TRS is murg, and each right-ground TRS is murg. For the concept of a finite path overlapping TRS (FPO-TRS) see [18]. For the concept of a layered transducing TRS (LT-TRS), see [19].

Let R be a rewrite system over Σ .

- (a) R is left-to-right minimal if for each rule $l \rightarrow r$ in R , $\rightarrow_{R-\{l \rightarrow r\}}^* \subset \rightarrow_R^*$.
- (b) R is left-to-right ground minimal if for each rule $l \rightarrow r$ in R , $\rightarrow_{R-\{l \rightarrow r\}}^* \cap (T_\Sigma \times T_\Sigma) \subset \rightarrow_R^* \cap (T_\Sigma \times T_\Sigma)$.

The set of all ground terms that are irreducible for a TRS R is denoted by $IRR(R)$.

Let R be a convergent TRS over Σ , and let $p \in T_\Sigma(X)$. It is well known that there exists exactly one term $t \in T_\Sigma(X)$ irreducible for R such that $p \rightarrow_R^* t$. We call t the R -normal form of p . We denote t by $p \downarrow_R$. Let $L \subseteq T_\Sigma$. The set of R -normal forms of the trees in the tree language L is denoted by $R(L)$. It should be clear that $R(L) = R^*(L) \cap IRR(R)$.

For the concept of a critical pair, see [1].

Let R and S be rewrite systems over the disjoint ranked alphabets Σ and Δ , respectively. Then the disjoint union $R \oplus S$ of R and S is the rewrite system $R \cup S$ over the ranked alphabet $\Sigma \cup \Delta$. Let $\mathbf{C}$ be a class of rewrite systems, let $\mathbf{C}$ be closed under disjoint union. A property $\mathcal{P}$ is modular for $\mathbf{C}$ if for any $R, S \in \mathbf{C}$ over disjoint ranked alphabets, $R \oplus S$ has the property $\mathcal{P}$ if and only if both R and S have the property $\mathcal{P}$.

2.3 Tree Languages

Let Σ be a ranked alphabet, a bottom-up tree automaton (bta) over Σ is a quadruple $\mathcal{A} = (\Sigma, A, R, A_f)$, where A is a finite set of states of rank 0, $\Sigma \cap A = \emptyset$, $A_f \subseteq A$ is the set of final states, R is a finite set of rules of the following two types:

- (i) $\delta(e_1, \dots, e_n) \rightarrow a$ with $n \geq 0$, $\delta \in \Sigma_n$, $a_1, \dots, a_n, a \in A$.
- (ii) $a \rightarrow a'$ with $a, a' \in A$ (λ -rules).

We consider R as a ground TRS over $\Sigma \cup A$. The tree language recognized by $\mathcal{A}$ is $L(\mathcal{A}) = \{t \in T_\Sigma \mid (\exists a \in A_f) t \rightarrow_R^* a\}$. A tree language L is recognizable if there exists a bta $\mathcal{A}$ such that $L(\mathcal{A}) = L$ (see [8]).

The bta $\mathcal{A} = (\Sigma, A, R, A_f)$ is deterministic if R has no λ -rules and R has no two rules with the same left-hand side.

Definition 2.1 Let Σ be a ranked alphabet, and let $L \subseteq T_\Sigma$ be a finite tree language. We define the *the fundamental bta* $\mathcal{A} = (\Sigma, A, R, A_f)$ of L as follows.

$$\begin{aligned}
A &= \{ \langle p \rangle \mid \text{there is } q \in L \text{ such that } p \in \text{sub}(q) \}. \\
R &= \{ \delta(\langle p_1 \rangle, \dots, \langle p_m \rangle) \rightarrow \langle p \rangle \mid \delta \in \Sigma_m, m \geq 0, \langle p_1 \rangle, \dots, \langle p_m \rangle, \langle p \rangle \in A, \text{ and} \\
&\quad p = \delta(p_1, \dots, p_m) \}. \\
A_f &= \{ \langle p \rangle \mid p \in L \}.
\end{aligned}$$

Lemma 2.2 *Let Σ be a ranked alphabet, and let $L \subseteq T_\Sigma$ be a finite tree language. Let $\mathcal{A} = (\Sigma, A, R, A_f)$ be the fundamental bta of L . Then $\mathcal{A}$ is deterministic, and $L(\mathcal{A}) = L$.*

Proof. By the definition of R , $\mathcal{A}$ is deterministic. It is not hard to see that for any $t \in \text{sub}(L)$ and $a \in A$, $t \rightarrow_A^* a$ if and only if $a = \langle t \rangle$. By the definition of A_f , $L(\mathcal{A}) = L$. □

3 TRSs Preserving Recognizability

Let Σ be a ranked alphabet, let R be a TRS over Σ , and let L be a tree language over Σ . Then $R_\Sigma^*(L) = \{p \mid q \rightarrow_R^* p \text{ for some } q \in L\}$ is the set of descendants of trees in L . When Σ is apparent from the context, we simply write $R^*(L)$ rather than $R_\Sigma^*(L)$.

For the concept of a $P\Sigma R$ -TRS, a $P\Sigma RF$ -TRS, a PR -TRS, a PRF -TRS, an $EP\Sigma R$ -TRS, an $EP\Sigma RF$ -TRS, an EPR -TRS, and an $EPRF$ -TRS, see the Introduction.

Theorem 3.1 *Each terminating TRS is an $EPRF$ -TRS.*

Proof. Let R be a terminating TRS over Σ and let $L \subseteq T_\Sigma$ be a finite language. Let $t \in T_\Sigma$ be arbitrary. We now show that $R^*(\{t\})$ is finite. On the contrary, assume that $R^*(\{t\})$ is infinite. Then t starts an infinite reduction sequence $t = t_0 \rightarrow_R t_1 \rightarrow_R t_2 \rightarrow_R t_3 \rightarrow_R \dots$ by König's lemma. Hence R is not terminating, which is a contradiction.

Thus $R^*(L)$ is finite and hence recognizable. We compute $R^*(L)$ as follows. Let $W = L$. While there is $q \in T_\Sigma - W$ such that $p \rightarrow_R q$ for some $p \in W$ we add q to W . Since $R^*(L)$ is finite, we stop. When we stop we have $W = R^*(L)$. Then we can construct a bta $\mathcal{C}$ over Σ such that $L(\mathcal{C}) = W$. □

Statement 3.2 *There is a left-linear monadic TRS R over a ranked alphabet Σ such that R is an $EP\Sigma RF$ -TRS, and that R is not a $P\Sigma R$ -TRS.*

Proof. Let $\Sigma = \Sigma_0 \cup \Sigma_2$, $\Sigma_0 = \{\sharp\}$, and $\Sigma_2 = \{f\}$. Let R over Σ consist of the rule $f(x_1, \sharp) \rightarrow f(x_1, x_1)$. Observe that $\Sigma = \text{sign}(R)$. We obtain by direct inspection that R is a left-linear monadic TRS R over Σ . Let $L \subseteq T_\Sigma$ be an arbitrary finite tree language. For any trees $p, q \in T_\Sigma$, if $p \rightarrow_R q$, then $\text{height}(p) = \text{height}(q)$. Hence $R^*(L)$ is finite. Thus $R^*(L)$ is recognizable. We compute $R^*(L)$ as follows. Let $W = L$. While there is $q \in T_\Sigma - W$ such that $p \rightarrow_R q$ for some $p \in W$ we add q to W . When we stop we have $W = R^*(L)$. We can construct a bta $\mathcal{C}$ over Σ such that $L(\mathcal{C}) = W$.

Then we define the n th left comb left_n for $n \geq 0$, as follows.

- (i) $\text{left}_0 = \sharp$, and
- (ii) for each $n \geq 0$, $\text{left}_{n+1} = f(\text{left}_n, x_{n+1})$.

Let

$$L = \{ \text{left}_n \mid n \geq 0 \}.$$

Then L is a recognizable tree language. Furthermore, for any $p \in R^*(L)$ and $s \in \text{sub}(p)$, if $s = f(t_1, t_2)$, then $t_2 = \#$ or $\text{height}(t_1) = \text{height}(t_2)$. For each $n \geq 0$, $f(\text{left}_n, \text{left}_n) \in R^*(L)$. Assume that $R^*(L)$ is a recognizable tree language. Similarly to the proof of the pumping lemma for recognizable tree languages, one can show the following. There are $0 \leq i < j$ such that $f(\text{left}_i, \text{left}_j) \in R^*(L)$. However, $\text{height}(\text{left}_i) = i < j = \text{height}(\text{left}_j)$. This contradicts our observation on $R^*(L)$. $\square$

Statement 3.3 *There is an LT-TRS R over a ranked alphabet Σ such that R is not a PΣRF-TRS.*

Proof. Let $\Sigma = \mathcal{F} \cup \mathcal{Q} = \Sigma_0 \cup \Sigma_1$, $\Sigma_0 = \{ \# \}$, $\Sigma_1 = \{ f, q \}$, $\mathcal{F} = \{ \#, f \}$, and $\mathcal{Q} = \{ q \}$. Let TRS R consist of the rules

$$\begin{aligned} f(q(x_1)) &\rightarrow q(f(x_1)), \\ \# &\rightarrow q(f(\#)). \end{aligned}$$

Then R is an LT-TRS. It is not hard to see that

$$R^*(\{ \# \}) \cap \{ f^k(q^m(\#)) \mid k, m \geq 0 \} = \{ f^k(q^k(\#)) \mid k \geq 0 \}.$$

Here $\{ f^k(q^m(\#)) \mid k, m \geq 0 \}$ is a recognizable tree language, and $\{ f^k(q^k(\#)) \mid k \geq 0 \}$ is not a recognizable tree language. The intersection of two recognizable tree languages is also a recognizable tree language. Hence $R^*(\{ \# \})$ is not a recognizable tree language. $\square$

Theorem 3.4 *There is a ranked alphabet Σ and a murg TRS R over Σ such that R is not a PΣRF-TRS.*

Proof. Let $\Sigma = \Sigma_0 \cup \Sigma_1 \cup \Sigma_2$, $\Sigma_0 = \{ \#, \$, \flat \}$, $\Sigma_1 = \{ f \}$, and $\Sigma_2 = \{ g, h \}$. Let the TRS R over Σ consist of the rules

$$\begin{aligned} \# &\rightarrow f(\#), \\ \# &\rightarrow \flat, \\ \$ &\rightarrow f(\$), \\ \$ &\rightarrow \flat, \\ g(x_1, x_1) &\rightarrow h(x_1, x_1), \end{aligned}$$

Consider a tree $h(t_1, t_2) \in R^*(g(\#, \$))$ where $t_1, t_2 \in T_\Sigma$. Then

$$g(\#, \$) \rightarrow_R^* g(f^k(\#), f^k(\$)) \rightarrow_R^* g(f^k(\flat), f^k(\flat)) \rightarrow_R h(f^k(\flat), f^k(\flat)) = h(t_1, t_2)$$

holds for some $k \geq 0$. Hence

$$R^*(\{ g(\#, \$) \}) \cap \{ h(t_1, t_2) \mid t_1, t_2 \in T_\Sigma \} = \{ h(f^k(\flat), f^k(\flat)) \mid k \geq 0 \}.$$

It is well known that the intersection of any two recognizable tree languages is a recognizable tree language. Observe that $\{ h(t_1, t_2) \mid t_1, t_2 \in T_\Sigma \}$ is a recognizable

tree language, and $\{h(f^k(b), f^k(b)) \mid k \geq 0\}$ is not a recognizable tree language. Thus $R^*(\{g(\$, \$)\})$ is not a recognizable tree language. $\square$

With an arbitrary Post Correspondence System (PCS) $\langle \mathbf{w}, \mathbf{z} \rangle$, Vágvolgyi [22] associated a ranked alphabet Σ , containing the distinguished nullary symbol $\# \in \Sigma_0$, and a TRS R over Σ . Vágvolgyi [22] showed the following results.

Statement 3.5 *If PCS $\langle \mathbf{w}, \mathbf{z} \rangle$ has a solution, then R is an $EP\Sigma R$ -TRS.*

Statement 3.6 *If PCS $\langle \mathbf{w}, \mathbf{z} \rangle$ has no solution, then $R^*(\{\#\})$ is not a recognizable tree language over Σ .*

Statement 3.6 implies the following statement.

Statement 3.7 *If PCS $\langle \mathbf{w}, \mathbf{z} \rangle$ has no solution, then R is not a $P\Sigma RF$ -TRS.*

The following result is a simple consequence of Statements 3.5 and 3.7.

Statement 3.8 *PCS $\langle \mathbf{w}, \mathbf{z} \rangle$ has a solution if and only if R is an $EP\Sigma RF$ -TRS if and only if TRS R is a $P\Sigma RF$ -TRS.*

Statement 3.8 implies the following result.

Proposition 3.9 *The following problem is undecidable:*

Instance: *A murg TRS R over a ranked alphabet Σ .*

Question: *Is R a $P\Sigma RF$ -TRS?*

We now recall the notion of a GSM-TRS, see [12].

Definition 3.10 Let R be a TRS over Σ . We say that R is a generalized semi-monadic TRS (GSM-TRS for short) if there is no rule $l \rightarrow r$ in R with $l \in X$ and the following holds. For any rules $l_1 \rightarrow r_1$ and $l_2 \rightarrow r_2$ in R , for any positions $\alpha \in POS(r_1)$ and $\beta \in POS(l_2)$, and for any supertree $l_3 \in T_\Sigma(X)$ of l_2/β with $var(l_3) \cap var(l_1) = \emptyset$, if

- (i) $\alpha = \lambda$ or $\beta = \lambda$,
- (ii) r_1/α and l_3 are unifiable, and
- (iii) σ is a most general unifier of r_1/α and l_3 ,

then

- (a) $l_2/\beta \in X$ or
- (b) for each $\gamma \in POS(l_3)$, if $l_2/\beta\gamma \in X$, then $\sigma(l_3/\gamma) \in X \cup T_\Sigma$.

Notice that Condition (a) implies that $l_3 \in X$.

Example 3.11 Let $\Sigma = \Sigma_0 \cup \Sigma_1 \cup \Sigma_3$, $\Sigma_0 = \{\#\}$, $\Sigma_1 = \{f\}$, and $\Sigma_3 = \{g\}$. Let the TRS R over Σ consist of the rule

$$g(x_1, x_2, \#) \rightarrow f(g(x_1, \#, x_1)).$$

We obtain by direct inspection that R is left-linear GSM-TRS.

The proof of the following result is straightforward.

Observation 3.12 *Each murg TRS is a GSM-TRS as well.*

Gylenizse and Vágvolgyi [12] observed that Fülöp's [5] undecidability results on deterministic top-down tree transducers simply imply the following.

Statement 3.13 [12] *Each of the following questions is undecidable for any convergent left-linear GSM-TRSs R and S over a ranked alphabet Ω , for any recognizable tree language $L \subseteq T_\Omega$ given by a tree automaton over Ω recognizing L , where $\Gamma \subseteq \Omega$ is the smallest ranked alphabet for which $R(L) \subseteq T_\Gamma$.*

- (i) *Is $R(L) \cap S(L)$ empty?*
- (ii) *Is $R(L) \cap S(L)$ infinite?*
- (iii) *Is $R(L) \cap S(L)$ recognizable?*
- (iv) *Is $T_\Gamma - R(L)$ empty?*
- (v) *Is $T_\Gamma - R(L)$ infinite?*
- (vi) *Is $T_\Gamma - R(L)$ recognizable?*
- (vii) *Is $R(L)$ recognizable?*
- (viii) *Is $R(L) = S(L)$?*
- (ix) *Is $R(L) \subseteq S(L)$?*

Theorem 3.14 *There is an FPO-TRS R such that R is not an PRF-TRS.*

Proof. Let $\Sigma = \Sigma_0 \cup \Sigma_1 \cup \Sigma_2$, $\Sigma_0 = \{ \$ \}$, $\Sigma_1 = \{ d, g \}$, $\Sigma_2 = \{ f \}$. Let the TRS R over Σ consist of the following rules.

$$\begin{aligned}
& \$ \rightarrow d(\$) \\
& g(d(x_1)) \rightarrow f(g(x_1), d(x_1)), \\
& g(\$) \rightarrow \$, \\
& f(\$, x_1) \rightarrow \$, \\
& f(\$, x_1) \rightarrow h(x_1, x_1).
\end{aligned}$$

By direct inspection of R , we get that R is an FPO-TRS. We now study the set $R^*(\{ \{ g(\$) \} \}) \cap \{ h(t_1, t_2) \mid t_1, t_2 \in T_\Sigma \}$. Assume that

$$g(\$) = u_0 \rightarrow_R u_2 \rightarrow_R u_3 \rightarrow_R \cdots \rightarrow_R u_{k-1} \rightarrow_R u_k = h(t_1, t_2)$$

for some $k \geq 1$ and $t_1, t_2 \in T_\Sigma$. We iterate application of the first and the second rules. We can change the order of applications of the first and second rules. Then we apply the third rule. Then we we apply the fourth rule finitely many times. We apply the fifth rule in the k th step, and hence $u_k = f(\$, t_1)$, and $t_1 = t_2$. Thus we obtain the following reduction sequence for some $n \geq 1$:

$$\begin{aligned}
& g(\$) \rightarrow_R g(d(\$)) \rightarrow_R g(d^2(\$)) \rightarrow_R \cdots \rightarrow_R g(d^n(\$)) \rightarrow_R \\
& f(g(d^{n-1}(\$), d^n(\$)) \rightarrow_R f(f(g(d^{n-2}(\$), d^{n-1}(\$)), d^n(\$)) \rightarrow_R \\
& f(f(f(g(d^{n-3}(\$), d^{n-2}(\$)), d^{n-1}(\$))), d^n(\$)) \rightarrow_R \cdots \rightarrow_R \\
& f(\dots f(f(f(g(\$), d(\$)), d^2(\$)), d^3(\$)), \dots, d^n(\$)) \rightarrow_R \\
& f(\dots f(f(f(g(\$), d(\$)), d^2(\$)), d^3(\$)), \dots, d^n(\$)) \rightarrow_R \\
& f(\dots f(f(f(\$, d(\$)), d^2(\$)), d^3(\$)), \dots, d^n(\$)) \rightarrow_R \\
& f(\dots f(f(\$, d^2(\$)), d^3(\$)), \dots, d^n(\$)) \rightarrow_R \\
& f(\dots f(\$, d^3(\$)), \dots, d^n(\$)) \rightarrow_R \cdots \rightarrow_R f(\$, d^n(\$)) \rightarrow_R h(d^n(\$), d^n(\$)).
\end{aligned}$$

In the light of the above reduction sequence, one can show that

$$R^*(\{\{g(\$)\}\}) \cap \{h(t_1, t_2) \mid t_1, t_2 \in T_\Sigma\} = \{h(d^n(\$), d^n(\$)) \mid n \geq 1\}.$$

Since $\{h(t_1, t_2) \mid t_1, t_2 \in T_\Sigma\}$ is a recognizable tree language, and $\{h(d^n(\$), d^n(\$)) \mid n \geq 1\}$ is not a recognizable tree language, we get that $R^*(\{g(\$)\})$ is not a recognizable tree language. $\square$

4 Main Results

We now show that each left-linear GSM-TRS is an EPRF-TRS.

Theorem 4.1 *Each left-linear GSM-TRS is an EPRF-TRS.*

Proof. Let R be a left-linear GSM-TRS over some ranked alphabet Σ . Moreover, let L be a finite tree language over Σ . Via a series of Lemmas we show that $R^*(L)$ is recognizable. To this end, we construct a bta $\mathcal{C}$ over Σ such that $L(\mathcal{C}) = R^*(L)$. Our construction is illustrated by an example in Section 4.

Let E be the set of all ground terms u over Σ such that there are rules $l_1 \rightarrow r_1$ and $l_2 \rightarrow r_2$ in R , and there are positions $\alpha \in POS(r_1)$ and $\beta \in POS(l_2)$, and there is a supertree $l_3 \in T_\Sigma(X) - X$ of l_2/β with $var(l_3) \cap var(l_1) = \emptyset$ such that

- (i) $\alpha = \lambda$ or $\beta = \lambda$,
- (ii) r_1/α and l_3 are unifiable, and
- (iii) σ is a most general unifier of r_1/α and l_3 , and
- (iv) there is a position $\gamma \in POS(l_3)$ such that $l_2/\beta\gamma \in X$ and $\sigma(l_3/\gamma) \in T_\Sigma$ and $u = \sigma(l_3/\gamma)$.

It should be clear that E is finite and is effectively constructable.

Moreover, without loss of generality we may assume that for each rule $l \rightarrow r$ in R , $l \in \bar{T}_\Sigma(X_n)$ for some $n \geq 0$. Let

$$D = sub(L) \cup \{p[e_1, \dots, e_n] \mid n \geq 0, p \in T_\Sigma(X_n), e_1, \dots, e_n \in sub(L \cup E), p \text{ is a subtree of the right-hand side } r \text{ of some rule } l \rightarrow r \text{ in } R\}.$$

Apparently, $sub(E) \subseteq D$. Hence $sub(L \cup E) \subseteq D$.

Let $\mathcal{A} = (\Sigma, A, S_A, A')$ be the fundamental bta of L . Recall that $\mathcal{A}$ is a deterministic bta over Σ such that $L(\mathcal{A}) = L$. Let $\mathcal{B} = (\Sigma, B, S_B, B')$ be the fundamental bta of D . Recall that $\mathcal{B}$ is a deterministic bta over Σ such that $L(\mathcal{B}) = D$. By the definition of D , we have $A \subseteq B$ and

$$S_A \subseteq S_B. \tag{1}$$

For each $i \geq 0$, consider the bta $\mathcal{C}_i = (\Sigma, B, S_i, A')$, where S_i is defined by recursion on i (for an example see Section 4). Let

$$S_0 = S_B. \tag{2}$$

Then $\mathcal{C}_0 = (\Sigma, B, S_B, A')$. Let us assume that $i \geq 1$ and we have defined the set S_{i-1} . Then we define S_i as follows.

(a) $S_{i-1} \subseteq S_i$.

(b) For any rule $l \rightarrow r$ in R with $n \geq 0$, $l \in \bar{T}_\Sigma(X_n)$, for all $e_1, \dots, e_n \in \text{sub}(L \cup E)$, if $l[\langle e_1 \rangle, \dots, \langle e_n \rangle] \rightarrow_{S_{i-1}}^* c$ for some $c \in B$, then we put the rule $\langle r[e_1, \dots, e_n] \rangle \rightarrow c$ in S_i .

By (1) and (2), we have

$$S_A \subseteq S_0. \quad (3)$$

It should be clear that there is an integer $M \geq 0$ such that $S_M = S_{M+1}$. Let M be the least integer such that $S_M = S_{M+1}$. Let $\mathcal{C} = \mathcal{C}_M$. Let $S = S_M$, and from now on we write $\mathcal{C} = (\Sigma, B, S, A')$, rather than $\mathcal{C}_M = (\Sigma, B, S_M, A')$.

Our aim is to show that $R^*(L) = L(\mathcal{C})$. To this end, first we show five preparatory lemmas, then the inclusion $L(\mathcal{C}) \subseteq R^*(L)$, then again five preparatory lemmas, and finally the inclusion $R^*(L) \subseteq L(\mathcal{C})$.

Lemma 4.2 $L = L(\mathcal{C}_0)$.

Proof. By Lemma 2.2, $L(\mathcal{A}) = L$. By the definition of $\mathcal{B}$ and $\mathcal{C}_0$, we have $L(\mathcal{C}_0) = L(\mathcal{A}) = L$. □

Lemma 4.3 For any $p \in T_\Sigma$ and $r \in T_\Sigma(X_n)$ with $n \geq 0$ and $\text{var}(r) = X_n$, if $p \rightarrow_{S_0}^* \langle r[e_1, \dots, e_n] \rangle$, then $p = r[e_1, \dots, e_n]$.

Proof. By direct inspection of the rules of S_0 . □

The following statement is a simple consequence of Lemma 4.3.

Lemma 4.4 For any $p \in T_\Sigma$, $r \in T_\Sigma(X_n)$, and $e_1, \dots, e_n \in \text{sub}(L \cup E)$, if $p \rightarrow_{S_0}^* \langle r[e_1, \dots, e_n] \rangle$, then $p = r[e_1, \dots, e_n]$.

Lemma 4.5 For any $i \geq 1$, $p \in T_\Sigma$, $q, t \in T_{\Sigma \cup B}$, $k \geq 1$, and $v_1, \dots, v_k \in T_{\Sigma \cup B}$, if

$$p = v_1 \xrightarrow{S_0} v_2 \xrightarrow{S_0} \dots \xrightarrow{S_0} v_k = q \xrightarrow{S_i} t, \quad (4)$$

and $\mathcal{C}_i$ applies an $(S_i - S_{i-1})$ -rule in the last step $q \rightarrow_{S_i} t$ of (4), then there exists an $s \in T_\Sigma$ such that

$$s \xrightarrow{R} p \text{ and } s \xrightarrow{S_{i-1}}^* t. \quad (5)$$

Proof. Let α be the position where $\mathcal{C}_i$ applies an $(S_i - S_{i-1})$ -rule

$$\langle r[e_1, \dots, e_n] \rangle \rightarrow c \text{ with } r \in T_\Sigma(X_n), n \geq 0, \text{ and } e_1, \dots, e_n \in \text{sub}(L \cup E)$$

in the last step $q \rightarrow_{S_i} t$ of (4). Then

$$q = u[\langle r[e_1, \dots, e_n] \rangle],$$

where $u \in \bar{T}_\Sigma(X_1)$, $u/\alpha = x_1$. By Lemma 4.4,

$$p = u[r[e_1, \dots, e_n]].$$

Finally, $t = u[c]$. By (b) of the definition of S_i , $i \geq 1$, there is a rule $l \rightarrow r$ in R with $l \in \bar{T}_\Sigma(X_n)$, $n \geq 0$ such that

$$l[\langle e_1 \rangle, \dots, \langle e_n \rangle] \xrightarrow[S_{i-1}]^* c.$$

Let

$$s = u[l[e_1, \dots, e_n]].$$

Then

$$s \xrightarrow[R]{} p$$

and

$$s = u[l[e_1, \dots, e_n]] \xrightarrow[S_0]^* u[l[\langle e_1 \rangle, \dots, \langle e_n \rangle]] \xrightarrow[S_{i-1}]^* u[c] = t.$$

Hence (5) holds. $\square$

Lemma 4.6 *For any $i \geq 0$, $p \in T_\Sigma$, and $q \in T_{\Sigma \cup B}$, if $p \rightarrow_{S_i}^* q$, then there is an $s \in T_\Sigma$ such that*

$$s \xrightarrow[R]{}^* p \text{ and } s \xrightarrow[S_0]{}^* q.$$

Proof. We proceed by induction on i . For $i = 0$ the statement is trivial. Let us suppose that $i \geq 1$ and that we have shown the statement for $1, 2, \dots, i-1$. Let

$$p \xrightarrow[S_i]{}^* q, \tag{6}$$

and let m be the number of $(S_i - S_{i-1})$ -rules applied by C_i along (6). We show by induction on m that

$$\text{there is } s \in T_\Sigma \text{ such that } s \xrightarrow[R]{}^* p \text{ and } s \xrightarrow[S_0]{}^* q. \tag{7}$$

If $m = 0$, then $p \rightarrow_{S_{i-1}}^* q$ and hence by the induction hypothesis on i , (7) holds.

Let us suppose that $m \geq 1$ and that for $0, 1, \dots, m-1$, we have shown (7). Let $p \rightarrow_{S_i}^* q$ where $\mathcal{C}$ applies m $(S_i - S_{i-1})$ -rules. Then there are integers n, k , $1 \leq k \leq n$, and there are trees $t_1, t_2, u_1, u_2, \dots, u_n \in T_{\Sigma \cup B}$ such that (I), (II), (III), and (IV) hold.

(I) $p = u_1 \rightarrow_{S_i} \dots \rightarrow_{S_i} u_k = t_1 \rightarrow_{S_i} u_{k+1} = t_2 \rightarrow_{S_i} \dots \rightarrow_{S_i} u_n = q$.

(II) along the reduction subsequence $p = u_1 \rightarrow_{S_i} \dots \rightarrow_{S_i} u_k = t_1$ of (I), $\mathcal{C}_i$ applies no $(S_i - S_{i-1})$ -rule.

(III) in the rewrite step $u_k \rightarrow_{S_i} u_{k+1}$ $\mathcal{C}_i$ applies an $(S_i - S_{i-1})$ -rule.

(IV) along the reduction subsequence $t_2 = u_{k+1} \rightarrow_{S_i} \dots \rightarrow_{S_i} u_n = q$ of (I), $\mathcal{C}_i$ applies $m-1$ $(S_i - S_{i-1})$ -rules.

By the induction hypothesis on i , there is a tree $s_1 \in T_\Sigma$ such that

$$s_1 \xrightarrow[R]{}^* p \text{ and } s_1 \xrightarrow[S_0]{}^* t_1. \tag{8}$$

Hence

$$s_1 \xrightarrow[S_0]{*} t_1 \xrightarrow[S_i]{} t_2.$$

By Lemma 4.5, there is a tree $s_2 \in T_\Sigma$ such that

$$s_2 \xrightarrow[R]{} s_1 \text{ and } s_2 \xrightarrow[S_{i-1}]{*} t_2. \quad (9)$$

Hence there is $j \geq 0$ and there are $w_1, \dots, w_j \in T_{\Sigma \cup B}$ such that

$$s_2 = w_1 \xrightarrow[S_{i-1}]{} w_2 \xrightarrow[S_{i-1}]{} \dots \xrightarrow[S_{i-1}]{} w_j = t_2 = u_{k+1} \xrightarrow[S_i]{} \dots \xrightarrow[S_i]{} u_n = q, \quad (10)$$

and along (10), $\mathcal{C}_i$ applies $m - 1$ $(S_i - S_{i-1})$ -rules. By the induction hypothesis on m , there is a tree $s_3 \in T_\Sigma$ such that

$$s_3 \xrightarrow[R]{} s_2 \text{ and } s_3 \xrightarrow[S_0]{*} q.$$

Hence by (8) and (9),

$$s_3 \xrightarrow[R]{} s_2 \xrightarrow[R]{} s_1 \xrightarrow[R]{} p.$$

Thus (7) holds. $\square$

Lemma 4.7 $L(\mathcal{C}) \subseteq R^*(L)$.

Proof. Let $p \in L(\mathcal{C})$. Then $p \rightarrow_S^* b$ for some $b \in A'$. Hence by Lemma 4.6, there is an $s \in T_\Sigma$ such that

$$s \xrightarrow[R]{} p \text{ and } s \xrightarrow[S_0]{*} b. \quad (11)$$

Hence $s \in L(\mathcal{C}_0)$. By Lemma 4.2, $s \in L$. Thus by (11), $p \in R^*(L)$. $\square$

Now we show the inclusion $R^*(L) \subseteq L(\mathcal{C})$. To this end, first we prove five lemmas.

Lemma 4.8 Let $l_1 \rightarrow r_1$ and $l_2 \rightarrow r_2$ be rules in R . Let $\alpha \in \text{POS}(r_1)$, where $r_1/\alpha \in T_\Sigma(X_j)$, $j \geq 0$. Let $\beta \in \text{POS}(l_2)$, where $l_2/\beta \in T_\Sigma(X) - X$, and let $s \in \bar{T}_\Sigma(X_k) - X$, $k \geq 1$, be a supertree of l_2/β . Let $\alpha = \lambda$ or $\beta = \lambda$. Let

$$(r_1/\alpha)[e_1, \dots, e_j] = s[z_1, \dots, z_k], \quad (12)$$

where $e_1, \dots, e_j \in \text{sub}(L \cup E)$, $z_1, \dots, z_k \in T_\Sigma$. Let $\gamma \in \text{POS}(s)$ be such that $l_2/\beta\gamma \in X$, and $s/\gamma = x_\nu$, for some $1 \leq \nu \leq k$. Then $z_\nu \in \text{sub}(L \cup E)$.

Proof. Let $l_1 \in T_\Sigma(X_m)$ for some $m \geq 0$. Let $l_3 = s[x_{m+1}, \dots, x_{m+k}]$. Then $l_3 \in T_\Sigma(\{x_{m+1}, \dots, x_{m+k}\})$ is a supertree of l_2/β , for each $m + 1 \leq i \leq m + k$, x_i appears exactly once in l_3 . Moreover, $\text{var}(l_1) \cap \text{var}(l_3) = \emptyset$, and by (12),

$$(r_1/\alpha)[e_1, \dots, e_j] = l_3[x_{m+1} \leftarrow z_1, \dots, x_{m+k} \leftarrow z_k]. \quad (13)$$

Let $\sigma_1 : X \rightarrow T_\Sigma(X)$ be a most general unifier of r_1/α and l_3 . By (13), there is a substitution $\sigma_2 : X \rightarrow T_\Sigma(X)$ such that

$$\sigma_2(\sigma_1(r_1/\alpha)) = (r_1/\alpha)[e_1, \dots, e_j] = l_3[x_{m+1} \leftarrow z_1, \dots, x_{m+k} \leftarrow z_k] = \sigma_2(\sigma_1(l_3)),$$

where $\sigma_2(\sigma_1(x_i)) = e_i$ for $1 \leq i \leq j$ and $\sigma_2(\sigma_1(x_{m+i})) = z_i$ for $1 \leq i \leq k$. Let $\gamma \in POS(s)$ be such that $l_2/\beta\gamma \in X$, and $s/\gamma = x_\nu$, for some $1 \leq \nu \leq k$. By Definition 3.10 and by the definition of E , $\sigma_1(x_{m+\nu}) \in X \cup sub(E)$. If $\sigma_1(x_{m+\nu}) \in X$, then $\sigma_2(\sigma_1(x_{m+\nu}))$ is a subtree of e_μ for some $\mu \in \{1, \dots, j\}$. Hence by the definition of $e_1, \dots, e_j$, $z_\nu = \sigma_2(\sigma_1(x_{m+\nu})) \in sub(L \cup E)$. If $\sigma_1(x_{m+\nu}) \in sub(E)$, then $z_\nu = \sigma_2(\sigma_1(x_{m+\nu})) = \sigma_1(x_{m+\nu}) \in sub(E)$. $\square$

Intuitively, the following lemma states that along a reduction sequence of S we can reverse the order of the consecutive application of a S_0 -rule at $\alpha \in N^*$ and the application of an $(S - S_0)$ -rule at $\beta \in N^*$ if α is not a prefix of β and β is not a prefix of α .

Lemma 4.9 *Let*

$$u_1 \xrightarrow{S} u_2 \xrightarrow{S} u_3$$

*be a reduction sequence of $\mathcal{C}$, where $u_1, u_2, u_3 \in T_{\Sigma \cup B}$. Let $\alpha \in POS(u_1)$, and $\beta \in POS(u_2)$ be such that $u_1 \xrightarrow{S} u_2$ applying a rule **rule₁** of S_0 at α , and that $u_2 \xrightarrow{S} u_3$ applying an $(S - S_0)$ -rule **rule₂** at β . If α is not a prefix of β and β is not a prefix of α , then there is a tree $v \in T_{\Sigma \cup B}$ such that $u_1 \xrightarrow{S} v$ applying **rule₂** at β , and $v \xrightarrow{S} u_3$ applying **rule₁** at α .*

Proof. Straightforward. $\square$

Lemma 4.10 *Let $i \geq 0$, $t \in \bar{T}_{\Sigma \cup B}(X_1)$, $\alpha \in POS(t)$, $t/\alpha = x_1$, $p \in D - sub(L)$, and $w \in sub(L)$. Let*

$$t[\langle p \rangle] = u_1 \xrightarrow{S_i} u_2 \xrightarrow{S_i} \dots \xrightarrow{S_i} u_n = \langle w \rangle \quad (14)$$

with $n \geq 1$, $u_1, \dots, u_n \in T_{\Sigma \cup B}$. Then along (14), $\mathcal{C}_i$ applies a rule in $S_i - S_0$ at some prefix β of α .

Proof. By direct inspection of the construction of the $\mathcal{C}_i$'s. $\square$

Lemma 4.11 *For any $n \geq 0$, $u \in \bar{T}_\Sigma(X_n)$, $v_1, \dots, v_n, v \in D$, $m \geq 1$, and $w_1, \dots, w_m \in T_{\Sigma \cup B}$, if*

$$u[\langle v_1 \rangle, \dots, \langle v_n \rangle] = w_1 \xrightarrow{S_0} w_2 \xrightarrow{S_0} \dots \xrightarrow{S_0} w_m = \langle v \rangle, \quad (15)$$

then $u[v_1, \dots, v_n] = v$.

Proof. We proceed by induction on $height(u)$. The basis $height(u) = 0$ of the induction is trivial. The induction step is a simple consequence of the definition of S_0 . $\square$

Lemma 4.12 *Let $t \in L(\mathcal{C})$, $m \geq 1$, $t_1, \dots, t_m \in T_{\Sigma \cup B}$, $b \in A'$, and let*

$$t = t_1 \xrightarrow{S} t_2 \xrightarrow{S} t_3 \xrightarrow{S} \dots \xrightarrow{S} t_m = b. \quad (16)$$

Let $l \rightarrow r$ be a rule in R , where $l \in \bar{T}_\Sigma(X_n)$ and $n \geq 1$. Moreover, let $1 \leq j \leq m$, and let

$$t_j/\alpha = l[\langle v_1 \rangle, \dots, \langle v_n \rangle], \quad (17)$$

where $n \geq 1$, $v_1, \dots, v_n \in D$, $\alpha \in POS(t_j)$. Let $\alpha_1, \dots, \alpha_n \in POS(l)$ be such that

$$l/\alpha_i = x_i \text{ for } 1 \leq i \leq n. \quad (18)$$

Consider the reduction subsequence

$$t_j \xrightarrow{S} t_{j+1} \xrightarrow{S} \dots \xrightarrow{S} t_m = b \quad (19)$$

of (16). If $\mathcal{C}$ does not apply any rules at the positions $\alpha\alpha_1, \dots, \alpha\alpha_n$ along (19), then $v_1, \dots, v_n \in \text{sub}(L \cup E)$.

Proof. Let $1 \leq i \leq n$, and let us assume that $v_i \in D - \text{sub}(L)$. By (17) and (18),

$$t_j/\alpha\alpha_i = \langle v_i \rangle. \quad (20)$$

By Lemma 4.10, $\mathcal{C}$ applies a rule in $S - S_0$ at some prefix of $\alpha\alpha_i$ along (19). Let $\beta \in POS(t_j)$ be the longest prefix of $\alpha\alpha_i$ such that $\mathcal{C}$ applies a rule **rule** in $S - S_0$ at β along (19). Then **rule** is of the form $\langle r_1[e_1, \dots, e_\kappa] \rangle \rightarrow c$, where $\kappa \geq 0$, $r_1 \in T_\Sigma(X_\kappa)$, $e_1, \dots, e_\kappa \in \text{sub}(L \cup E)$, and there is a rule $l_1 \rightarrow r_1$ in R . Moreover there exists ξ , $j < \xi \leq m$, such that

$$t_j/\beta \xrightarrow{S}^* t_{j+1}/\beta \xrightarrow{S}^* \dots \xrightarrow{S}^* t_\xi/\beta = \langle r_1[e_1, \dots, e_\kappa] \rangle,$$

where for each π , $j \leq \pi \leq \xi - 1$, $t_\pi/\beta = t_{\pi+1}/\beta$ or $t_\pi/\beta \xrightarrow{S} t_{\pi+1}/\beta$. We lose no generality by assuming that

$$t_j/\beta \xrightarrow{S} t_{j+1}/\beta \xrightarrow{S} \dots \xrightarrow{S} t_\xi/\beta = \langle r_1[e_1, \dots, e_\kappa] \rangle. \quad (21)$$

By Lemma 4.9 we may assume that there exists ν , $j \leq \nu \leq \xi$ such that

(a) along the reduction subsequence

$$t_j/\beta \xrightarrow{S} \dots \xrightarrow{S} t_\nu/\beta \quad (22)$$

of (21) no rule is applied at any prefix of $\alpha\alpha_i$, that

(b) along (22) each application of a rule of S_0 at some $\delta \in N^*$ is followed somewhere later by an application of an $S - S_0$ -rule of S at a prefix ϵ of δ , and that

(c) along the reduction subsequence

$$t_\nu/\beta \xrightarrow{S} \dots \xrightarrow{S} t_\xi/\beta = \langle r_1[e_1, \dots, e_\kappa] \rangle$$




Figure 1: Case 1.

of (21), S applies only rules of S_0 .

Then

$$t_\nu/\beta = s[\langle z_1 \rangle, \dots, \langle z_k \rangle] \quad (23)$$

for some $k \geq 1$, $s \in \bar{T}_\Sigma(X_k)$, and $z_1, \dots, z_k \in D$. By (23), (c) of the definition of ν , and Lemma 4.11,

$$s[z_1, \dots, z_k] = r_1[e_1, \dots, e_\kappa]. \quad (24)$$

The word α is a prefix of β or β is a prefix of α . Hence we can distinguish two cases.

Case 1 α is a prefix of β , see Figure 4. In this case,

$$\beta = \alpha\gamma \quad (25)$$

for some $\gamma \in N^*$, and hence t_ν/β is a subtree of t_ν/α . Now by (17), the definition of ν , and (23),

$$s \text{ is a supertree of } l/\gamma. \quad (26)$$

Let ω be the prefix of $\alpha\alpha_i$ with $\text{length}(\omega) = \text{length}(\alpha\alpha_i) - 1$. Observe that $\mathcal{C}$ applies a rule of S_0 at the position ω along (19). Hence

$$s \notin X. \quad (27)$$

We define $\delta \in N^*$ be by the equation $\gamma\delta = \alpha_i$. Then

$$\beta\delta = \alpha\alpha_i, \quad (28)$$

and by (a) of the definition of ν ,

$$\delta \in \text{POS}(s), \delta \in \text{POS}(l/\gamma), \text{ and } (l/\gamma)/\delta = x_i. \quad (29)$$




Figure 2: Case 2.

By (28) and by (a) of the definition of ν ,

$$\beta\delta \in POS(t_\nu).$$

By (20), (28), (a) of the definition of ν , and (23),

$$\langle v_i \rangle = (t_j/\beta)/\delta = (t_\nu/\beta)/\delta = s[\langle z_1 \rangle, \dots, \langle z_k \rangle]/\delta = \langle z_\mu \rangle \quad (30)$$

for some $1 \leq \mu \leq k$. As R is a GSM-TRS, by (26), (27), (29), (24), and Lemma 4.8, $z_\mu \in sub(L \cup E)$. By (30), $v_i = z_\mu$. Thus $v_i \in sub(L \cup E)$.

Case 2 β is a prefix of α , see Figure 5. In this case

$$\alpha = \beta\gamma \quad (31)$$

for some $\gamma \in N^*$, and hence t_j/α is a subtree of t_j/β . Now by (17), the definition of ν , and (23),

$$s/\gamma \text{ is a supertree of } l. \quad (32)$$

Moreover, by (a) of the definition of ν ,

$$\alpha_i \in POS(s/\gamma), \quad l/\alpha_i \in X, \quad \text{and} \quad (s/\gamma)/\alpha_i \in X. \quad (33)$$

Let ω be the prefix of $\alpha\alpha_i$ with $length(\omega) = length(\alpha\alpha_i) - 1$. Observe that $\mathcal{C}$ applies a rule of S_0 at the position ω along (19). Hence

$$s/\gamma \notin X. \quad (34)$$

By (31) and by (a) of the definition of ν ,

$$\beta\gamma\alpha_i = \alpha\alpha_i \in POS(t_\nu). \quad (35)$$

Then by (20), (35), (a) of the definition of ν , and (23),

$$\langle v_i \rangle = (t_j/\beta)/\gamma\alpha_i = (t_\nu/\beta)/\gamma\alpha_i = s[\langle z_1 \rangle, \dots, \langle z_k \rangle]/\gamma\alpha_i = \langle z_\mu \rangle \quad (36)$$

for some $1 \leq \mu \leq k$. By (24),

$$(s/\gamma)[z_1, \dots, z_k] = s[z_1, \dots, z_k]/\gamma = r_1[e_1, \dots, e_\kappa]/\gamma. \quad (37)$$

As R is a GSM-TRS, by (32), (34), (33), (36), (37), and Lemma 4.8, $z_\mu \in \text{sub}(L \cup E)$. By (36), $v_i = z_\mu$. Thus $v_i \in \text{sub}(L \cup E)$. $\square$

Lemma 4.13 $R^*(L) \subseteq L(\mathcal{C})$.

Proof. By (3), $L \subseteq L(\mathcal{C}_0)$. As $S_{i-1} \subseteq S_i$ for $i \geq 1$, we have $L \subseteq L(\mathcal{C}_i)$ for $i \geq 0$. Hence $L \subseteq L(\mathcal{C})$. Thus it is sufficient to show that for each $t \in L(\mathcal{C})$, if $t \rightarrow_R t'$, then $t' \in L(\mathcal{C})$. To this end, let us suppose that $t \rightarrow_R t'$, applying the rule $l \rightarrow r$ in R at $\alpha \in \text{POS}(t)$. Here $l \in \bar{T}_\Sigma(X_n)$ for some $n \geq 0$. Let $\alpha_1, \dots, \alpha_n \in \text{POS}(l)$ be such that

$$l/\alpha_i = x_i \text{ for } 1 \leq i \leq n.$$

Then

$$t = s[l[u_1, \dots, u_n]],$$

where $s \in \bar{T}_\Sigma(X_1)$, $\alpha \in \text{POS}(s)$, $s/\alpha = x_1$, and $u_1, \dots, u_n \in T_\Sigma$. Moreover,

$$t' = t[\alpha \leftarrow r[u_1, \dots, u_n]] = s[r[u_1, \dots, u_n]].$$

As $t \in L(\mathcal{C})$, there is a reduction sequence

$$t = t_1 \xrightarrow{S} t_2 \xrightarrow{S} t_3 \xrightarrow{S} \dots \xrightarrow{S} t_m = b, \quad (38)$$

where $m \geq 1$, $b \in A'$, $t_1, \dots, t_m \in T_{\Sigma \cup B}$, and there are integers j, k with $1 \leq j \leq k \leq m$ such that

- (i) $t_j = s[l[\langle v_1 \rangle, \dots, \langle v_n \rangle]]$, where $v_i \in D$ and $u_i \rightarrow_S^* \langle v_i \rangle$ for $1 \leq i \leq n$,
- (ii) $t_k = s[c_0]$, for some $c_0 \in A$, where $l[\langle v_1 \rangle, \dots, \langle v_n \rangle] \rightarrow_S^* c_0$, and that
- (iii) along the reduction subsequence $t_j \rightarrow_S t_{j+1} \rightarrow_S \dots \rightarrow_S t_k$ of (38), $\mathcal{C}$ does not apply any rules at the positions $\alpha\alpha_1, \dots, \alpha\alpha_n$. By Lemma 4.12, $v_1, \dots, v_n \in \text{sub}(L \cup E)$. Hence by Condition (b) in the definition of S_i , $i \geq 1$, and by the definition of $\mathcal{C}$, the rule $r[\langle v_1 \rangle, \dots, \langle v_n \rangle] \rightarrow c_0$ is in S . Thus we get

$$t' = s[r[u_1, \dots, u_n]] \xrightarrow{S}^* s[r[\langle v_1 \rangle, \dots, \langle v_n \rangle]] \xrightarrow{S} s[c_0] \xrightarrow{S}^* b.$$

As $b \in A'$, we have $t' \in L(\mathcal{C})$. $\square$

By Lemma 4.7 and Lemma 4.13, we get that $R^*(L) = L(\mathcal{C})$. $\square$

Lemma 3.2 and Theorem 4.1 imply the following result.

Theorem 4.14 *There is a left-linear monadic TRS R over a ranked alphabet Σ such that R is an EPRF-TRS and that R is not a $P\Sigma R$ -TRS.*

5 An Example

We illustrate the construction of $\mathcal{C}_j$, $j \geq 0$, appearing in the previous section by an example. Let $\Sigma = \Sigma_0 \cup \Sigma_1 \cup \Sigma_3$, $\Sigma_0 = \{\#\}$, $\Sigma_1 = \{f\}$, $\Sigma_3 = \{g\}$. Let the TRS R over Σ consist of the following two rules.

$$\begin{aligned} f(f(g(x_1, \#, \#))) &\rightarrow f(f(x_1)), \\ g(x_1, x_2, \#) &\rightarrow f(g(x_1, \#, x_1)). \end{aligned}$$

By direct inspection we obtain that R is a left-linear GSM-TRS. Here $E = \{\#\}$. Let $L = \{g(\#, \#, \#)\}$. Then $sub(L \cup E) = \{\#, g(\#, \#, \#)\}$. It is not hard to see that

$$R^*(L) = \{f^n(g(\#, \#, \#)) \mid n \geq 0\} \cup \{f^n(\#) \mid n \geq 2\}.$$

By direct inspection we obtain that the set of subterms of the right-hand sides of the rules of R is

$$\{x_1, f(x_1), f(f(x_1)), \#, g(x_1, \#, x_1), f(g(x_1, \#, x_1))\}.$$

Then

$$\begin{aligned} D &= \{\#, f(\#), g(\#, \#, \#), f(f(\#)), f(g(\#, \#, \#)), f(f(g(\#, \#, \#))), g(g(\#, \#, \#), \#, g(\#, \#, \#)), \\ &\quad f(g(g(\#, \#, \#), \#, g(\#, \#, \#)))\}. \\ \mathcal{C}_0 &= (\Sigma, B, S_0, \{\langle g(\#, \#, \#) \rangle\}), \text{ where} \\ B &= \{\langle \# \rangle, \langle g(\#, \#, \#) \rangle, \langle f(\#) \rangle, \langle f(g(\#, \#, \#)) \rangle, \langle f(f(\#)) \rangle, \langle f(f(g(\#, \#, \#))) \rangle, \\ &\quad \langle g(g(\#, \#, \#), \#, g(\#, \#, \#)) \rangle, \langle f(g(g(\#, \#, \#), \#, g(\#, \#, \#))) \rangle\}. \end{aligned}$$

Furthermore, S_0 consists of the following eight rules.

$$\begin{aligned} \# &\rightarrow \langle \# \rangle, \\ g(\langle \# \rangle, \langle \# \rangle, \langle \# \rangle) &\rightarrow \langle g(\#, \#, \#) \rangle, \\ f(\langle \# \rangle) &\rightarrow \langle f(\#) \rangle, \\ f(\langle g(\#, \#, \#) \rangle) &\rightarrow \langle f(g(\#, \#, \#)) \rangle, \\ f(f(\langle \# \rangle)) &\rightarrow \langle f(f(\#)) \rangle, \\ f(\langle f(g(\#, \#, \#)) \rangle) &\rightarrow \langle f(f(g(\#, \#, \#))) \rangle, \\ g(\langle g(\#, \#, \#) \rangle, \langle \# \rangle, \langle g(\#, \#, \#) \rangle) &\rightarrow \langle g(g(\#, \#, \#), \#, g(\#, \#, \#)) \rangle, \\ f(\langle g(g(\#, \#, \#), \#, g(\#, \#, \#)) \rangle) &\rightarrow \langle f(g(g(\#, \#, \#), \#, g(\#, \#, \#))) \rangle. \end{aligned}$$

$\mathcal{C}_1 = (\Sigma, B, S_1, \{\langle g(\#, \#, \#) \rangle\})$, where S_1 contains all rules of S_0 and the following rules.

$$\begin{aligned} \langle f(f(\#)) \rangle &\rightarrow \langle f(f(g(\#, \#, \#))) \rangle, \\ \langle f(g(\#, \#, \#)) \rangle &\rightarrow \langle g(\#, \#, \#) \rangle, \end{aligned}$$

$\mathcal{C}_2 = (\Sigma, B, S_2, \{\langle g(\#, \#, \#) \rangle\})$, where S_2 contains all rules of S_1 and the following two rules.

$$\begin{aligned} \langle f(f(\#)) \rangle &\rightarrow \langle f(g(\#, \#, \#)) \rangle, \\ \langle f(f(\#)) \rangle &\rightarrow \langle g(\#, \#, \#) \rangle. \end{aligned}$$

The bta $\mathcal{C}_3 = (\Sigma, B, S_3, \{\langle g(\#, \#, \#) \rangle\})$ is equal to $\mathcal{C}_2$. By direct inspection we obtain that the states

$$\begin{aligned} &\langle f(f(g(\#, \#, \#))) \rangle, \langle g(g(\#, \#, \#), \#, g(\#, \#, \#)) \rangle, \\ &\langle f(g(g(\#, \#, \#), \#, g(\#, \#, \#))) \rangle \end{aligned}$$

are superfluous as the final state $\langle g(\#, \#, \#) \rangle$ cannot be reached from any of them. Hence we drop all of them and also omit all rules in which they appear. In this way we obtain the bta $\mathcal{B}_1 = (\Sigma, B_1, Q_1, \{ \langle g(\#, \#, \#) \rangle \})$, where

$B_1 = \{ \langle \# \rangle, \langle g(\#, \#, \#) \rangle, \langle f(\#) \rangle, \langle f(g(\#, \#, \#)) \rangle, \langle f(f(\#)) \rangle \}$ and Q_1 consists of the following rules.

$\# \rightarrow \langle \# \rangle,$
 $g(\langle \# \rangle, \langle \# \rangle, \langle \# \rangle) \rightarrow \langle g(\#, \#, \#) \rangle,$
 $f(\langle \# \rangle) \rightarrow \langle f(\#) \rangle,$
 $f(f(\langle \# \rangle)) \rightarrow \langle f(f(\#)) \rangle,$
 $\langle f(g(\#, \#, \#)) \rangle \rightarrow \langle g(\#, \#, \#) \rangle,$
 $\langle f(f(\#)) \rangle \rightarrow \langle f(g(\#, \#, \#)) \rangle,$
 $\langle f(f(\#)) \rangle \rightarrow \langle g(\#, \#, \#) \rangle,$

We obtain the bta $\mathcal{B}_2 = (\Sigma, B_2, Q_2, \{ \langle g(\#, \#, \#) \rangle \})$ from $\mathcal{B}_1$ by eliminating the lambda rules. Here $B_2 = B_1$ and Q_2 consists of the following rules.

$\# \rightarrow \langle \# \rangle,$
 $g(\langle \# \rangle, \langle \# \rangle, \langle \# \rangle) \rightarrow \langle g(\#, \#, \#) \rangle,$
 $f(\langle \# \rangle) \rightarrow \langle f(\#) \rangle,$
 $f(\langle f(\#) \rangle) \rightarrow \langle f(f(\#)) \rangle.$
 $f(\langle f(\#) \rangle) \rightarrow \langle f(g(\#, \#, \#)) \rangle,$
 $f(f(\langle \# \rangle)) \rightarrow \langle g(\#, \#, \#) \rangle,$
 $f(\langle g(\#, \#, \#) \rangle) \rightarrow \langle g(\#, \#, \#) \rangle.$

By direct inspection we obtain that the states $\langle f(f(\#)) \rangle$ and $\langle f(g(\#, \#, \#)) \rangle$ are superfluous as the final state $\langle g(\#, \#, \#) \rangle$ cannot be reached from any of them. Hence we drop all of them and also omit all rules in which they appear.

In this way we obtain the bta $\mathcal{B}_3 = (\Sigma, B_3, Q_3, \{ \langle g(\#, \#, \#) \rangle \})$.

Here $B_3 = \{ \langle \# \rangle, \langle g(\#, \#, \#) \rangle, \langle f(\#) \rangle, \}$ and

Q_3 consists of the following five rules.

$\# \rightarrow \langle \# \rangle,$
 $g(\langle \# \rangle, \langle \# \rangle, \langle \# \rangle) \rightarrow \langle g(\#, \#, \#) \rangle,$
 $f(\langle \# \rangle) \rightarrow \langle f(\#) \rangle,$
 $f(\langle f(\#) \rangle) \rightarrow \langle g(\#, \#, \#) \rangle,$
 $f(\langle g(\#, \#, \#) \rangle) \rightarrow \langle g(\#, \#, \#) \rangle.$

Then $L(\mathcal{C}_3) = L(\mathcal{B}_3)$. We obtain by direct inspection that $L(\mathcal{B}_3) = R^*(L)$.

6 PRF-TRSs

We show various decidability and undecidability results on PRF-TRSs and EPRF-TRSs. We show that reachability, joinability, and local confluence are decidable for EPRF-TRSs.

Theorem 6.1 *There is a ranked alphabet Σ and there is a linear $EP\Sigma RF$ -TRS R such that R is not a PRF-TRS.*

Proof. Let $\Sigma = \Sigma_1 \cup \Sigma_0$, $\Sigma_1 = \{f, g\}$, $\Sigma_0 = \{\#\}$. Let R consist of the following five rules.

$$\begin{aligned} f(g(x_1) \rightarrow f(f(g(g(x_1))))), \\ f(\#) \rightarrow \#, \\ g(\#) \rightarrow \#, \\ \# \rightarrow f(\#), \\ \# \rightarrow g(\#). \end{aligned}$$

It should be clear that for each tree $t \in T_\Sigma$, $t \rightarrow_R^* \#$ and $\# \rightarrow_R^* t$. Hence for each nonempty tree language $L \subseteq T_\Sigma$, $R^*(L) = T_\Sigma$. Thus R is an EP Σ RF-TRS.

Let $\Delta = \Sigma \cup \{h\}$, where $h \in \Delta_1$. Then $R^*(\{f(g(h(\#)))\}) = \{f^n(g^n(h(t))) \mid n \geq 0, t \in T_\Sigma\}$ is not recognizable. $\square$

Theorem 6.2 *Let R be any TRS over $\text{sign}(R)$, and let $\Sigma = \{f, \#\} \cup \text{sign}(R)$, where $f \in \Sigma_2 - \text{sign}(R)$ and $\# \in \Sigma_0 - \text{sign}(R)$. Then R is a P Σ RF-TRS if and only if R is a PRF-TRS.*

Proof. ($\Leftarrow$) Trivial.

($\Rightarrow$) Let Γ be an arbitrary ranked alphabet with $\text{sign}(R) \subseteq \Gamma$. To each symbol $g \in \Gamma_k - \text{sign}(R)$, $k \geq 0$, we assign a tree $t_g \in T_\Sigma(X_k)$. To this end, we number the symbols in $\Gamma - \text{sign}(R)$ from 1 to $|\Gamma - \text{sign}(R)|$.

Then we define the n th right comb right_n for $n \geq 0$, as follows.

- (i) $\text{right}_0 = \#$,
- (ii) for each $n \geq 0$, $\text{right}_{n+1} = f(\#, \text{right}_n)$.

For the definition of the n th left comb left_n , see the proof of Statement 3.2. To any symbol $g \in \Gamma_k - \text{sign}(R)$, $k \geq 0$, with number l , we assign the tree $t_g = f(\text{left}_k, \text{right}_l)$.

Consider the TRS

$$S = \{g(x_1, \dots, x_k) \rightarrow t_g \mid k \geq 0, g \in \Gamma_k - \text{sign}(R)\}.$$

It should be clear that S is a convergent TRS. It is not hard to show the following two statements.

Claim 6.3 *For any $r, s \in T_\Gamma$,*

$$r \xrightarrow[R]{} s \text{ if and only if } r \downarrow_S \xrightarrow[R]{} s \downarrow_S.$$

Claim 6.4 *A tree language L over $\text{sign}(R)$ is finite if and only if the tree language $S(L)$ over Σ is finite. A tree language L over $\text{sign}(R)$ is recognizable if and only if the tree language $S(L)$ over Σ is recognizable.*

Let L be any finite tree language over $\text{sign}(R)$. By Claim 6.4, $S(L)$ is a finite tree language over Σ . By Claim 6.3, $S(R_{\text{sign}(R)}^*(L)) = R_\Sigma^*(S(L))$. By Claim 6.4, $R_{\text{sign}(R)}^*(L)$ is recognizable if and only if $R_\Sigma^*(S(L))$ is recognizable. Hence if R is a P Σ RF-TRS, then R is a PRF-TRS. As Γ is an arbitrary ranked alphabet with $\text{sign}(R) \subseteq \Gamma$, R is a PRF-TRS. $\square$

The proof of the following result is similar to the proof of Theorem 6.2.

Theorem 6.5 *Let R be a TRS over $\text{sign}(R)$, and let $\Sigma = \{f, \sharp\} \cup \text{sign}(R)$, where $f \in \Sigma_2 - \text{sign}(R)$ and $\sharp \in \Sigma_0 - \text{sign}(R)$. R is an EPΣRF-TRS if and only if R is an EPRF-TRS.*

Consequence 6.6 *Let R be a TRS over Σ such that there is a symbol $f \in \Sigma_2 - \text{sign}(R)$ and there is a constant $\sharp \in \Sigma_0 - \text{sign}(R)$. Then R is a PRF-TRS if and only if R is a PΣRF-TRS. Moreover, R is an EPRF-TRS if and only if R is an EPΣRF-TRS.*

We now show that reachability is decidable for EPRF-TRSs.

Theorem 6.7 *Let R be an EPRF-TRS over Σ and let $p, q \in T_\Sigma(X)$. Then it is decidable whether $p \rightarrow_R^* q$.*

Proof. Let $m \geq 0$ be such that $\text{var}(p) \subseteq X_m$, $\text{var}(q) \subseteq X_m$. Let us introduce new constant symbols $Z = \{z_1, \dots, z_m\}$ with $Z \cap \Sigma = \emptyset$. For each $t \in T_\Sigma(X_m)$, we define $t_z \in T_{\Sigma \cup Z}$ as $t_z = t[z_1, \dots, z_m]$. By direct inspection we obtain that for all $u, v \in T_\Sigma(X)$,

$$u \xrightarrow[R]{} v \text{ if and only if } u_z \xrightarrow[R]{} v_z,$$

hence

$$u \xrightarrow[R]^* v \text{ if and only if } u_z \xrightarrow[R]^* v_z.$$

Consider the singleton set $\{p_z\}$. As R is an EPRF-TRS, $R_{\Sigma \cup Z}^*(\{p_z\})$ is a recognizable tree language over $\Sigma \cup Z$, and we can construct a bta over $\Sigma \cup Z$ which recognize $R_{\Sigma \cup Z}^*(\{p_z\})$. Hence we can decide whether $q_z \in R_{\Sigma \cup Z}^*(\{p_z\})$, see [8]. Clearly, $q_z \in R_{\Sigma \cup Z}^*(\{p_z\})$ if and only if $p \rightarrow_R^* q$. □

We now show that joinability is decidable for EPRF-TRSs.

Theorem 6.8 *Let R be an EPRF-TRS over Σ , and let $p, q \in T_\Sigma(X)$. Then it is decidable whether there is a tree $r \in T_\Sigma(X)$ such that $p \rightarrow_R^* r$ and $q \rightarrow_R^* r$.*

Proof. For each $t \in T_\Sigma(X_m)$, we define $t_z \in T_{\Sigma \cup Z}$ as in the proof of Theorem 6.7.

Claim 6.9 *For any $p, q \in T_\Sigma(X)$, there is a tree $r \in T_\Sigma(X)$ such that $p \rightarrow_R^* r$ and $q \rightarrow_R^* r$ if and only if $R_{\Sigma \cup Z}^*(\{p_z\}) \cap R_{\Sigma \cup Z}^*(\{q_z\}) = \emptyset$.*

Proof. Assume that $R_{\Sigma \cup Z}^*(\{p_z\}) \cap R_{\Sigma \cup Z}^*(\{q_z\}) = \emptyset$. Then there is a tree $s \in R_{\Sigma \cup Z}^*(\{p_z\}) \cap R_{\Sigma \cup Z}^*(\{q_z\})$. We define r from s by replacing each occurrence of z_i by x_i for $1 \leq i \leq m$. Then $p \rightarrow_R^* r$ and $q \rightarrow_R^* r$.

Assume that there is a tree $r \in T_\Sigma(X)$ such that $p \rightarrow_R^* r$ and $q \rightarrow_R^* r$. Hence $r_z \in R_{\Sigma \cup Z}^*(\{p_z\})$ and $r_z \in R_{\Sigma \cup Z}^*(\{q_z\})$. Thus $R_{\Sigma \cup Z}^*(\{p_z\}) \cap R_{\Sigma \cup Z}^*(\{q_z\}) \neq \emptyset$. □

As R is an EPRF-TRS, $R_{\Sigma \cup Z}^*(\{p_z\})$ and $R_{\Sigma \cup Z}^*(\{q_z\})$ are recognizable, and we can construct two btas over $\Sigma \cup Z$ which recognize $R_{\Sigma \cup Z}^*(\{p_z\})$ and $R_{\Sigma \cup Z}^*(\{q_z\})$, respectively. Hence we can decide whether $R_{\Sigma \cup Z}^*(\{p_z\}) \cap R_{\Sigma \cup Z}^*(\{q_z\}) = \emptyset$, see [8].

By Claim 6.9, if $R_{\Sigma \cup Z}^*(\{p_z\}) \cap R_{\Sigma \cup Z}^*(\{q_z\}) \neq \emptyset$, then there is a tree $r \in T_\Sigma(X)$ such that $p \rightarrow_R^* r$ and $q \rightarrow_R^* r$. Otherwise, there is no tree $r \in T_\Sigma(X)$ such that $p \rightarrow_R^* r$ and $q \rightarrow_R^* r$.

□

Theorem 6.10 *Let R be a confluent EPRF-TRS over Σ , and let $p, q \in T_\Sigma(X)$. Then it is decidable whether $p \leftrightarrow_R^* q$.*

Proof. $p \leftrightarrow_R^* q$ if and only if there is a tree $r \in T_\Sigma(X)$ such that $p \rightarrow_R^* r$ and $q \rightarrow_R^* r$. By Theorem 6.8, we can decide whether there is a tree $r \in T_\Sigma(X)$ such that $p \rightarrow_R^* r$ and $q \rightarrow_R^* r$.

□

We now show that local confluence is decidable for EPRF-TRSs.

Theorem 6.11 *Let R be an EPRF-TRS over Σ . Then it is decidable whether R is locally confluent.*

Proof. It is well known that R is locally confluent if and only if for every critical pair (v_1, v_2) of R there exists a tree $v \in T_\Sigma(X)$ such that $v_1 \rightarrow_R^* v$ and $v_2 \rightarrow_R^* v$, see [1]. Furthermore, all critical pairs of R are variants of finitely many critical pairs of R . Hence it is sufficient to inspect finitely many critical pairs. Thus the theorem follows from Theorem 6.8.

□

Theorem 6.12 *Let R be an EPRF-TRS and S be a TRS over Σ . Then it is decidable whether $\rightarrow_S^* \subseteq \rightarrow_R^*$.*

Proof. Let $m \geq 0$ be such that for all variables x_i occurring on the left-hand side of some rule in S , $x_i \in X_m$, that is, $i \leq m$. From now on, for each $t \in T_\Sigma(X_m)$, we define $t_z \in T_{\Sigma \cup Z}$ as in the proof of Theorem 6.7.

Claim 6.13 $\rightarrow_S^* \subseteq \rightarrow_R^*$ if and only if for each rule $l \rightarrow r$ in S , $r_z \in R_{\Sigma \cup Z}^*(\{l_z\})$.

Proof. ($\Rightarrow$) Let $l \rightarrow r$ be an arbitrary rule in S . Clearly, $l \rightarrow_R^* r$. Thus $r_z \in R_{\Sigma \cup Z}^*(\{l_z\})$.

($\Leftarrow$) Let us suppose that $t_1, t_2 \in T_\Sigma(X)$, and that $t_1 \rightarrow_S t_2$ applying the rule $l \rightarrow r$. As $r_z \in R_{\Sigma \cup Z}^*(\{l_z\})$, $l_z \rightarrow_R^* r_z$ holds. Hence $l \rightarrow_R^* r$ implying that $t_1 \rightarrow_R^* t_2$ as well.

□

Let $l \rightarrow r$ be an arbitrary rule in S . We can construct a bta over $\Sigma \cup Z$ recognizing the singleton set $\{l_z\}$. As R is an EPRF-TRS, $R_{\Sigma \cup Z}^*(\{l_z\})$ is recognizable, and we can construct a bta over $\Sigma \cup Z$ recognizing $R_{\Sigma \cup Z}^*(\{l_z\})$. Hence we can decide whether $r_z \in R_{\Sigma \cup Z}^*(\{l_z\})$. Thus by Claim 6.13, we can decide whether $\rightarrow_S^* \subseteq \rightarrow_R^*$.

□

Consequence 6.14 *Let R and S be EPRF-TRS over Σ . Then it is decidable which one of the following four mutually excluding conditions holds.*

- (i) $\rightarrow_R^* \subset \rightarrow_S^*$,
- (ii) $\rightarrow_S^* \subset \rightarrow_R^*$,
- (iii) $\rightarrow_R^* = \rightarrow_S^*$,
- (iv) $\rightarrow_R^* \bowtie \rightarrow_S^*$,

where “ $\bowtie$ ” stands for the incomparability relationship.

Observation 6.15 *If one omits a rule from a left-linear GSM-TRS, then the resulting rewrite system still remains a left-linear GSM-TRS.*

One can easily show the following result applying Theorem 4.1, Consequence 6.14, and Observation 6.15.

Consequence 6.16 *For a left-linear GSM-TRS R , it is decidable whether R is left-to-right minimal.*

Consequence 6.14 also implies the following.

Consequence 6.17 *Let R and S be TRSs over Σ such that $R \cup R^{-1}$ and $S \cup S^{-1}$ are EPRF-TRSs. Then it is decidable which one of the following four mutually excluding conditions holds.*

- (i) $\leftrightarrow_R^* \subset \leftrightarrow_S^*$,
- (ii) $\leftrightarrow_S^* \subset \leftrightarrow_R^*$,
- (iii) $\leftrightarrow_R^* = \leftrightarrow_S^*$,
- (iv) $\leftrightarrow_R^* \bowtie \leftrightarrow_S^*$.

Theorem 6.18 *Let R be an EPRF-TRS and S be a TRS over a ranked alphabet Σ . Let $g \in \Sigma - (\text{sign}(R) \cup \Sigma_0)$. Let $\sharp \in \Sigma_0$ be irreducible for R . Then it is decidable whether $\rightarrow_S^* \cap (T_\Sigma \times T_\Sigma) \subseteq \rightarrow_R^* \cap (T_\Sigma \times T_\Sigma)$.*

Proof. We assume that $g \in \Sigma_1$. One can easily modify the proof of this case when proving the more general case $g \in \Sigma_k$, $k \geq 1$. For each $t \in T_\Sigma(X)$, we define $t_g \in T_\Sigma$ from t by substituting $g^i(\sharp)$ for all occurrences of the variable x_i for $i \geq 1$.

Claim 6.19 $\rightarrow_S^* \cap (T_\Sigma \times T_\Sigma) \subseteq \rightarrow_R^* \cap (T_\Sigma \times T_\Sigma)$ if and only if for each rule $l \rightarrow r$ in S , $r_g \in R_1^*(\{l_g\})$.

Proof. ($\Rightarrow$) Let $l \rightarrow r$ be an arbitrary rule in S . Clearly, $l_g \rightarrow_S r_g$. Thus by our assumption $l_g \rightarrow_R^* r_g$.

($\Leftarrow$) Let us suppose that $t_1, t_2 \in T_\Sigma$, and that $t_1 \rightarrow_S t_2$ applying the rule $l \rightarrow r$. As $r_g \in R_1^*(\{l_g\})$, $l_g \rightarrow_R^* r_g$ holds. Hence $l \rightarrow_R^* r$ implying that $t_1 \rightarrow_R^* t_2$ as well. $\square$

For each rule $l \rightarrow r$ in S , the tree language $\{l_g\}$ is recognizable, and we can construct a bta over Σ recognizing $\{l_g\}$. As R is an EPRF-TRS, $R^*(\{l_g\})$ is also recognizable, and we can construct a bta over Σ recognizing $R^*(\{l_g\})$. Hence for each rule $l \rightarrow r$ in S , we can decide whether or not $r_g \in R^*(\{l_g\})$. Thus by Claim 6.19, we can decide whether $\rightarrow_S^* \cap (T_\Sigma \times T_\Sigma) \subseteq \rightarrow_R^* \cap (T_\Sigma \times T_\Sigma)$. $\square$

Consequence 6.20 *Let R and S be EPRF-TRSs over Σ . Moreover, let $g_1, g_2 \in \Sigma - \Sigma_0$ be such that for each $i \in \{1, 2\}$, g_i does not occur on the left-hand side of any rule in R_i . Let $\sharp_1, \sharp_2 \in \Sigma_0$ be such that for each $i \in \{1, 2\}$, $\sharp_i$ is irreducible for R_i . Then it is decidable which one of the following four mutually excluding conditions holds.*

- (i) $\rightarrow_R^* \cap (T_\Sigma \times T_\Sigma) \subset \rightarrow_S^* \cap (T_\Sigma \times T_\Sigma)$,
- (ii) $\rightarrow_S^* \cap (T_\Sigma \times T_\Sigma) \subset \rightarrow_R^* \cap (T_\Sigma \times T_\Sigma)$,
- (iii) $\rightarrow_R^* \cap (T_\Sigma \times T_\Sigma) = \rightarrow_S^* \cap (T_\Sigma \times T_\Sigma)$,
- (iv) $\rightarrow_R^* \cap (T_\Sigma \times T_\Sigma) \bowtie \rightarrow_S^* \cap (T_\Sigma \times T_\Sigma)$.

One can easily show the following result applying Theorem 4.1, Observation 6.15, and Consequence 6.20.

Consequence 6.21 *Let R be a left-linear GSM-TRS over Σ . Moreover, let $g \in \Sigma - \Sigma_0$ such that g does not occur on the left-hand side of any rule in R , and let $\sharp \in \Sigma_0$ be irreducible for R . Then it is decidable whether R is left-to-right ground minimal.*

By Statement 3.13 and Theorem 4.1 we have the following.

Theorem 6.22 *Each of the following questions is undecidable for any convergent left-linear EPR-TRSs R and S over a ranked alphabet Ω , for any recognizable tree language $L \subseteq T_\Omega$ given by a tree automaton over Ω recognizing L , where $\Gamma \subseteq \Omega$ is the smallest ranked alphabet for which $R(L) \subseteq T_\Gamma$.*

- (i) *Is $R(L) \cap S(L)$ empty?*
- (ii) *Is $R(L) \cap S(L)$ infinite?*
- (iii) *Is $R(L) \cap S(L)$ recognizable?*
- (iv) *Is $T_\Gamma - R(L)$ empty?*
- (v) *Is $T_\Gamma - R(L)$ infinite?*
- (vi) *Is $T_\Gamma - R(L)$ recognizable?*
- (vii) *Is $R(L)$ recognizable?*
- (viii) *Is $R(L) = S(L)$?*
- (ix) *Is $R(L) \subseteq S(L)$?*

Lemma 6.23 *Let R and S be linear collapse-free rewrite systems over the disjoint ranked alphabets Σ and Δ , respectively. Let Γ be a ranked alphabet with $\Sigma \cup \Delta \subseteq \Gamma$. Consider R and S as rewrite systems over Γ . Then*

- (i) $\rightarrow_S \circ \rightarrow_R \subseteq \rightarrow_R \cup (\rightarrow_R \circ \rightarrow_S)$, and
- (ii) $\rightarrow_{R \cup S}^* = \rightarrow_R^* \circ \rightarrow_S^*$.

Proof. The proof of (i) is straightforward. Condition (ii) is a simple consequence of (i). $\square$

Theorem 6.24 *Let R be a linear collapse-free EPRF-TRS and S be a linear collapse-free EPR-TRS over the disjoint ranked alphabets $\text{sign}(R)$ and $\text{sign}(S)$, respectively. Then $R \oplus S$ is a linear collapse-free EPR-TRS.*

Proof. Apparently, $R \oplus S$ is a linear collapse-free TRS. Let L be a recognizable tree language over some ranked alphabet Γ , where $\text{sign}(R) \cup \text{sign}(S) \subseteq \Gamma$. By Lemma 6.23, $(R \oplus S)_\Gamma^*(L) = S_\Gamma^*(R_\Gamma^*(L))$. As R is an EPRF-TRS, $R_\Gamma^*(L)$ is recognizable. Moreover, since S preserves recognizability, $S_\Gamma^*(R_\Gamma^*(L))$ is also recognizable. $\square$
The proof of the following result is similar to the proof of Theorem 6.24.

Theorem 6.25 *Let R be a linear collapse-free PRF-TRS and S be a linear collapse-free PR-TRS over the disjoint ranked alphabets Σ and Δ , respectively. Then $R \oplus S$ is a PR-TRS.*

Theorem 6.26 *Let R and S be TRSs over the disjoint ranked alphabets Σ and Δ , respectively, such that any left-hand side in $R \oplus S$ differs from a variable. If $R \oplus S$ is an EPRF-TRS, then R and S are also EPRF-TRSs.*

Proof. Let L be a finite recognizable tree language over some ranked alphabet Γ , where $\Sigma \subseteq \Gamma$. It is sufficient to show that $R_\Gamma^*(L)$ is recognizable. Without loss of generality we may rename the symbols of Γ such that $\Gamma \cap \Delta = \emptyset$. Thus $R_\Gamma^*(L) = (R \oplus S)_{\Gamma \cup \Delta}^*(L)$. Since $\Sigma \cup \Delta \subseteq \Gamma \cup \Delta$ and $R \oplus S$ is an EPRF-TRS, we get that $R_\Gamma^*(L)$ is recognizable and we can effectively construct a bta recognizing $R_\Gamma^*(L)$. $\square$

7 Conclusion and Open Problems

We showed that each left-linear GSM-TRS is an EPRF-TRS. We showed that reachability, joinability, and local confluence are decidable for EPRF-TRSs. We showed that the following problem is undecidable:

Instance: A murg TRS R over a ranked alphabet Σ .

Question: Is R a PΣRF-TRS?

Our results give rise to several open problems.

- Generalize the notion of a left-linear GSM-TRS such that the obtained TRS is still an EPRF-TRS.

- Show the following conjecture. Let R be a right-linear TRS over $\text{sign}(R)$, and let $\Sigma = \{g, \sharp\} \cup \text{sign}(R)$, where $g \in \Sigma_1 - \text{sign}(R)$ and $\sharp \in \Sigma_0 - \text{sign}(R)$. Then R is an EPΣRF-TRS if and only if R is an EPRF-TRS. Show the corresponding conjectures when R is left-linear or R is linear.

- Show that a string rewrite system R over the alphabet $\text{alph}(R)$ of R preserves $\text{alph}(R)$ -recognizability of finite string languages if and only if R preserves recognizability of finite string languages. Show that it is not decidable for a string rewrite system R whether R preserves $\text{alph}(R)$ -recognizability of finite string languages, and whether R preserves recognizability of finite string languages. Hence it is not decidable for a linear TRS R whether R is a PΣRF-TRS and whether R is a PRF-TRS.

- Show that the property preserving recognizability of finite tree languages and the property effectively preserving recognizability of finite tree languages are modular for the class of all left-linear collapse-free TRSs, for the class of all right-linear

collapse-free TRSs, for the class of all linear collapse-free TRSs, and for the class of all collapse-free TRSs.

References

- [1] F. Baader and T. Nipkow Term Rewriting and All That, Cambridge University Press, Cambridge, United Kingdom, 1998.
- [2] J.L. Coquidé, M. Dauchet, R. Gilleron, and S. Vágvölgyi, Bottom-up tree push-down automata: classification and connection with rewrite systems, *Theoretical Computer Science* **127** (1994) 69-98.
- [3] I. Durand, G. Sénizergues, Bottom-Up Rewriting Is Inverse Recognizability Preserving, in: F. Baader ed., Term Rewriting and Applications, 18th International Conference, RTA 2007, Paris, France, June 26-28, 2007, Proceedings. *Lecture Notes in Computer Science* **4533** Springer Verlag 2007, 107-121.
- [4] I. Durand, G. Sénizergues, Bottom-up rewriting for words and terms, arXiv:0903.2554v1 [cs.FL], <http://arxiv.org/abs/0903.2554>
- [5] Z. Fülöp, Undecidable properties of deterministic top-down tree transducers, *Theoretical Computer Science* **134** (1994) 311-328.
- [6] Z. Fülöp and P. Gyenizse, On injectivity of deterministic top-down tree transducers, *Information Processing Letters* **48** (1993) 183-188.
- [7] Z. Fülöp, E. Jurvanen, M. Steinby, and S. Vágvölgyi, On one-pass term rewriting, Mathematical Foundations of Computer Science, 1998 (Brno), Lecture Notes in Computer Science, **1450**, Springer Publishing Company, Berlin, 1998 248-256; see also in Acta Cybernetica, **14** (1999) 83-98.
- [8] F. Gécseg and M. Steinby, *Tree Automata* (Akadémiai Kiadó, Budapest, 1984).
- [9] F. Gécseg and M. Steinby, Tree Languages, in: G. Rozenberg and A. Salomaa eds., Handbook of Formal Language Theory to be issued by Springer Verlag in 1996.
- [10] R. Gilleron, Decision problems for term rewriting systems and recognizable tree languages, Proc. STACS'91, *Lecture Notes in Computer Science* **480**, (1991) 148-159.
- [11] R. Gilleron and S. Tison, Regular tree languages and rewrite systems, *Fundamenta Informaticae* **24** (1995) 157-175.
- [12] P. Gyenizse, S. Vágvölgyi: Linear Generalized Semi-Monadic Rewrite Systems Effectively Preserve Recognizability. *Theoretical Computer Science* 194(1-2): 87-122 (1998).

- [13] P. Gyvenizse, S. Vágvolgyi, A property of left-linear rewrite systems preserving recognizability, *Theoretical Computer Science* **242** (2000) 477-498.
- [14] F. Jacquemard, M. Rusinowitch: Closure of Hedge-Automata Languages by Hedge Rewriting, in: Andrei Voronkov ed., *Rewriting Techniques and Applications*, 19th International Conference, RTA 2008, Proceedings. Lecture Notes in Computer Science **5117** Springer (2008) 157-171.
- [15] Nagaya, T., Toyama, Y.: Decidability for left-linear growing term rewriting systems. *Information and Computation* 178(2), 499-514 (2002).
- [16] F. Otto, Some undecidability results concerning the property of preserving regularity, *Theoretical Computer Science* **207** (1998) 43-72.
- [17] K. Salomaa, Deterministic tree pushdown automata and monadic tree rewriting systems, *Journal of Computer and System Sciences* **37** (1988) 367-394.
- [18] T. Takai, Y. Kaji, and H. Seki, Right-Linear Finite Path Overlapping Term Rewriting Systems Effectively Preserve Recognizability, in: L. Bachmair ed., *Rewriting Techniques and Applications*, Lecture Notes in Computer Science **1833** (2000) 246-260.
- [19] T. Takai, H. Seki, Y. Fujinaka, and Y. Kaji, Layered Transducing Term Rewriting System and Its Recognizability Preserving Property, *IEICE Transactions on Information and Systems*, **E86-D** 285-295, 2003.
- [20] S. Vágvolgyi, Right-linear half-monadic term rewrite systems, *Theoretical Computer Science* **309** (2003) 195-211.
- [21] S. Vágvolgyi, Descendants of a recognizable tree language for sets of linear monadic term rewrite rules, *Information Processing Letters*, **99** (2006) 111-118.
- [22] S. Vágvolgyi, Murg term rewrite systems, *Information Processing Letters*, **108** (2008), 264-272.